## Math 3100 Assignment 8 Continuity and Differentiation

Due at 1:00 pm on Friday the 2nd of November 2018

- 1. Let  $f:[0,1] \to \mathbb{R}$  be continuous with f(0) = f(1). Show that there must exist  $x, y \in [0,1]$  satisfying |x-y| = 1/2 and f(x) = f(y).
- 2. Give an example of each of the following, or provide a short argument for why the request is impossible.
  - (a) A continuous function defined on [0,1] with range (0,1).
  - (b) A continuous function defined on (0,1) with range [0,1].
  - (c) A continuous function defined on (0,1] with range (0,1).
- 3. Suppose f is a continuous function, f(1) = -4, f(-2) = 3,  $\lim_{x \to -\infty} f(x) = 2$  and  $\lim_{x \to \infty} f(x) = -1$ . Prove that there exist  $c, d \in \mathbb{R}$  so that  $f(c) \le f(x) \le f(d)$  for all  $x \in \mathbb{R}$ .
- 4. Exactly one of the following requests is impossible. Decide which it is, and provide examples for the other three. In each case, lets assume that the functions are defined on all of  $\mathbb{R}$ .
  - (a) Function f and g not differentiable at  $x_0 = 0$ , but where fg is differentiable at  $x_0 = 0$ .
  - (b) A function f not differentiable at  $x_0 = 0$  and a function g differentiable at  $x_0 = 0$  where fg is differentiable at  $x_0 = 0$ .
  - (c) A function f not differentiable at  $x_0 = 0$  and a function g differentiable at  $x_0 = 0$  where f + g is differentiable at  $x_0 = 0$ .
  - (d) A function f differentiable at  $x_0 = 0$ , but not differentiable at any other point.
- 5. Use the definition of the derivative to find  $f'(x_0)$  for all  $x_0 \in \mathbb{R}$  if:

(a) 
$$f(x) = \sqrt{x^2 + 1}$$
 (b)  $f(x) = \frac{1}{x^2 + 1}$ 

- 6. (a) Let  $f(x) = \begin{cases} x^2, & \text{if } x \ge 0 \\ 0, & \text{if } x < 0 \end{cases}$ 
  - i. Compute f'(x) for  $x \neq 0$ .
  - ii. Use the definition of the derivative to find f'(0).
  - iii. Is f' continuous at 0? Give your reasoning.
  - iv. Does f''(0) exist? Give your reasoning.
  - (b) Let  $g(x) = \begin{cases} x^3 \sin \frac{1}{x^2}, & x \neq 0 \\ 0, & x = 0 \end{cases}$ .
    - i. Compute q'(x) for  $x \neq 0$ .
    - ii. Use the definition of the derivative to find g'(0).
    - iii. Is g' continuous at 0? Give your reasoning.
    - iv. Does g''(0) exist? Give your reasoning.
- 7. Prove that if g is differentiable at  $x_0$ , and  $g(x_0) \neq 0$ , then 1/g is differentiable at  $x_0$  and

$$\left(\frac{1}{g}\right)'(x_0) = \frac{-g'(x_0)}{g(x_0)^2}.$$

- 8. (a) Suppose f is continuous on [a, b], twice differentiable on (a, b), and  $f''(x) \neq 0$  for all  $x \in (a, b)$ . Prove carefully that f has at most 2 distinct zeros in [a, b].

  Hint: Use Rolle's Theorem
  - (b) Prove that the function  $f(x) = x^2 \sin x$  has precisely two roots.