

Sample Exam 2 – Version 1

No calculators. Show your work. Give full explanations. Good luck!

1. (15 points)

- (a) Carefully state what it means to say that $\sum_{n=1}^{\infty} a_n$ converges to A and prove that if this indeed the case, then $\sum_{n=1}^{\infty} (10a_n)$ converges to $10A$.
- (b) Prove that if $b_n > 0$ for all $n \in \mathbb{N}$ and $\sum_{n=1}^{\infty} b_n$ converges, then $\sum_{n=1}^{\infty} b_n^2$ also converges.
- (c) Prove that if a series converges absolutely, then it is convergent.

2. (15 points)

- (a) Determine if the following series are absolutely convergent, conditionally convergent, or divergent. Justify your answers.

$$(i) \quad \sum_{n=1}^{\infty} \frac{(-1)^n n}{n^2 + 1} \qquad (ii) \quad \sum_{n=1}^{\infty} \frac{\log n}{n^{3/2}}$$

- (b) Use the “Cauchy Condensation Test” to determine the convergence or divergence of

$$\sum_{n=2}^{\infty} \frac{1}{n \log n}$$

- (c) Find all $x \in \mathbb{R}$ for which $\sum_{n=1}^{\infty} \frac{(-2)^n x^{2n}}{n}$ converges.

3. (20 points) Let $f : \mathbb{R} \rightarrow \mathbb{R}$.

- (a) Carefully state the ε - δ definition of what it means for f to be *continuous* at x_0 and conclude that if f is continuous at x_0 with $f(x_0) = 2$, then there exists $\delta > 0$ such that $f(x) \geq 1$ whenever $|x - x_0| < \delta$.
- (b) Use the definition from part (a) to prove that $f(x) = \frac{1}{x}$ is continuous at $x_0 = 1$.
- (c) Prove that f is continuous at x_0 if and only if $\lim_{n \rightarrow \infty} f(x_n) = f(x_0)$ for all sequences with $\lim_{n \rightarrow \infty} x_n = x_0$.

Math 3100 - Sample Exam 2 (Version 1) - SOLUTIONS

1. (a) We say that $\sum_{n=1}^{\infty} a_n$ converges to A if

$$\lim_{n \rightarrow \infty} \underbrace{(a_1 + a_2 + \dots + a_n)}_{=: s_n \text{ ("n-th partial sum")}} \text{ exists \& equals } A.$$

Claim

If $\sum_{n=1}^{\infty} a_n$ converges to A , then $\sum_{n=1}^{\infty} (10a_n)$ converges to $10A$.

Proof

We know that $\lim_{n \rightarrow \infty} (a_1 + \dots + a_n) = A$. It follows that

$$\lim_{n \rightarrow \infty} \underbrace{(10a_1 + 10a_2 + \dots + 10a_n)}_{\text{"constant times limit law for sequences."}} = 10 \lim_{n \rightarrow \infty} (a_1 + a_2 + \dots + a_n) = 10A.$$

This is the n -th partial sum for the series $\sum_{n=1}^{\infty} 10a_n$. \square

(b) Claim: If $b_n > 0$ for all $n \in \mathbb{N}$ and $\sum_{n=1}^{\infty} b_n$ converges, then $\sum_{n=1}^{\infty} b_n^2$ converges also.

Proof. Since $\sum_{n=1}^{\infty} b_n$ converges we know that $\lim_{n \rightarrow \infty} b_n = 0$ and

hence that $\exists N$ such that $0 < b_n \leq 1$ for all $n > N$.

$$\Rightarrow 0 < b_n^2 \leq b_n \text{ for all } n > N.$$

It thus follows by "Direct Comparison" that $\sum_{n=1}^{\infty} b_n^2$ converges. \square

(c) Claim

If $\sum_{n=1}^{\infty} |a_n|$ converges, then $\sum_{n=1}^{\infty} a_n$ converges also.

Proof 1 (Using the "Cauchy Criterion for Series").

Recall: Cauchy Criterion for Series

$\sum_{n=1}^{\infty} b_n$ converges $\Leftrightarrow \forall \varepsilon > 0 \exists N$ such that if $n > m > N$, then $\left| \sum_{k=m+1}^n b_k \right| < \varepsilon$.

Let $\varepsilon > 0$. Since $\sum_{n=1}^{\infty} |a_n|$ converges we know that $\exists N$ such that if $n > m > N$, then $\left| \sum_{k=m+1}^n |a_k| \right| = \sum_{k=m+1}^n |a_k| < \varepsilon$.

Since $\left| \sum_{k=m+1}^n a_k \right| = |a_{m+1} + a_{m+2} + \dots + a_n|$

\triangle -inequality $\nearrow \leq |a_{m+1}| + |a_{m+2}| + \dots + |a_n| = \sum_{k=m+1}^n |a_k|$

it follows that if $n > m > N$, then $\left| \sum_{k=m+1}^n a_k \right| < \varepsilon$ and hence

by direction \Leftarrow of the Cauchy Criterion for Series that

$\sum_{n=1}^{\infty} a_n$ converges. □

Proof 2 (Using "Direct Comparison")

Since $-|a_n| \leq a_n \leq |a_n|$ for all $n \in \mathbb{N}$

$\Rightarrow 0 \leq a_n + |a_n| \leq 2|a_n|$ for all $n \in \mathbb{N}$.

Since $\sum_{n=1}^{\infty} (2|a_n|)$ converges it follows by "Direct Comparison" that $\sum_{n=1}^{\infty} (a_n + |a_n|)$ also converges.

Since $a_n = (a_n + |a_n|) - |a_n|$ and both

$$\sum_{n=1}^{\infty} (a_n + |a_n|) \text{ \& } \sum_{n=1}^{\infty} |a_n| \text{ converge}$$

it follows that $\sum_{n=1}^{\infty} a_n$ also converges. (by "Difference Limit Law",

□.

2. (a) (i) $\sum_{n=1}^{\infty} \frac{(-1)^n n}{n^2+1}$ CONVERGES CONDITIONALLY.

Since $\sum_{n=1}^{\infty} \frac{n}{n^2+1}$ diverges. (since $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges and

But $\sum_{n=1}^{\infty} \frac{(-1)^n n}{n^2+1}$ converges $\lim_{n \rightarrow \infty} \frac{\frac{n}{n^2+1}}{\frac{1}{n}} = \lim_{n \rightarrow \infty} \frac{n^2}{n^2+1} = 1$ by "Limit Comparison".

[By Alt. Series Test, since $\frac{n}{n^2+1} \rightarrow 0$.]

(ii) $\sum_{n=1}^{\infty} \frac{\log n}{n^{3/2}}$ CONVERGES ABSOLUTELY.

Since $\log n \leq n^{1/4}$ for all sufficiently large n implies.

$$\frac{\log n}{n^{3/2}} \leq \frac{n^{1/4}}{n^{3/2}} = \frac{1}{n^{5/4}} \text{ "eventually" \& } \sum_{n=1}^{\infty} \frac{1}{n^{5/4}} \text{ converges.}$$

(b) Since $\frac{1}{n \log n} \gg 0$ and

$$\sum_{k=1}^{\infty} 2^k \frac{1}{2^k \log(2^k)} = \sum_{k=1}^{\infty} \frac{1}{(\log 2) k} \quad \underline{\text{diverges}} \quad (\text{right?})$$

it follows from the "Cauchy Condensation Test" that

$$\sum_{n=2}^{\infty} \frac{1}{n \log n} \quad \text{diverges also.}$$

(c) Claim $\sum_{n=1}^{\infty} \frac{(-2)^n x^{2n}}{n}$ converges $\Leftrightarrow x \in [-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}]$.

Proof Let $a_n = \frac{(-2)^n x^{2n}}{n}$.

$$\text{Since } \left| \frac{a_{n+1}}{a_n} \right| = \left| \frac{(-2)^{n+1} x^{2n+2}}{n+1} \cdot \frac{n}{(-2)^n x^{2n}} \right|$$

$$= (2) \left(\frac{n}{n+1} \right) |x|^2 \rightarrow 2|x|^2$$

it follows from the "Ratio Test" that $\sum a_n$ converges absolutely if $|x| < \frac{1}{\sqrt{2}}$ and diverges if $|x| > \frac{1}{\sqrt{2}}$.

If $|x| = \frac{1}{\sqrt{2}}$, then $\sum_{n=1}^{\infty} a_n = \sum_{n=1}^{\infty} \frac{(-1)^n}{n}$ which converges

by the Alt. Series test since $\frac{1}{n} \gg 0$.

□

3. (a) A function $f: \mathbb{R} \rightarrow \mathbb{R}$ is continuous at a point $x_0 \in \mathbb{R}$ if $\forall \varepsilon > 0 \exists \delta > 0$ such that if $|x - x_0| < \delta$, then $|f(x) - f(x_0)| < \varepsilon$.

Claim If $f: \mathbb{R} \rightarrow \mathbb{R}$ is continuous at x_0 and $f(x_0) = 2$, then $\exists \delta > 0$ such that if $|x - x_0| < \delta$, then $f(x) > 1$.

Proof

Since f is conts at x_0 & $f(x_0) = 2$ it follows (with $\varepsilon = 1$) that $\exists \delta > 0$ such that if $|x - x_0| < \delta$, then

$$|f(x) - f(x_0)| < 1$$

$$\Updownarrow \leftarrow \text{since } f(x_0) = 2$$

$$1 < f(x) < 3$$

□

(b) Claim $f(x) = \frac{1}{x}$ is continuous at $x_0 = 1$.

Proof let $\varepsilon > 0$ and set $\delta = \min \left\{ \frac{1}{2}, \frac{\varepsilon}{2} \right\}$.

If $|x - 1| < \delta$ it follows that

$$\left| \frac{1}{x} - \frac{1}{1} \right| = \frac{|x - 1|}{|x|} < 2|x - 1| < 2\left(\frac{\varepsilon}{2}\right) = \varepsilon.$$

□

$$\begin{aligned} &\leftarrow \text{since } |x - 1| < \frac{1}{2} \\ &\Rightarrow |x| > \frac{1}{2} \Leftrightarrow \frac{1}{|x|} < 2 \end{aligned}$$

(c) Claim

$f: \mathbb{R} \rightarrow \mathbb{R}$ conts at $x_0 \iff \lim_{n \rightarrow \infty} f(x_n) = f(x_0)$ for all seqs $x_n \rightarrow x_0$.

Proof

(\Rightarrow) Let $\{x_n\}$ be a sequence with $\lim_{n \rightarrow \infty} x_n = x_0$ and $\varepsilon > 0$.

Since f conts at x_0 we know $\exists \delta > 0$ such that if

$$|x - x_0| < \delta \text{ then } |f(x) - f(x_0)| < \varepsilon.$$

Since $x_n \rightarrow x_0$ we know $\exists N$ such that if $n > N$ then

$$|x_n - x_0| < \delta$$

and hence $|f(x_n) - f(x_0)| < \varepsilon$ as required.

(\Leftarrow) [Contrapositive]

Suppose f is not continuous at x_0 . This means $\exists \varepsilon_0 > 0$ such that for every $\delta > 0$ there exists $x \in \mathbb{R}$ with $|x - x_0| < \delta$, but $|f(x) - f(x_0)| \geq \varepsilon_0$.

In particular, $\exists x_1 \in \mathbb{R}$ with $|x_1 - x_0| < 1$ and $|f(x_1) - f(x_0)| \geq \varepsilon_0$
 $\exists x_2 \in \mathbb{R}$ with $|x_2 - x_0| < \frac{1}{2}$ and $|f(x_2) - f(x_0)| \geq \varepsilon_0$

\vdots

In fact, $\forall n \in \mathbb{N} \exists x_n \in \mathbb{R}$ with $|x_n - x_0| < \frac{1}{n}$ but $|f(x_n) - f(x_0)| \geq \varepsilon_0$

Clearly, $\lim_{n \rightarrow \infty} x_n = x_0$ (by "Baby Squeeze"), but $f(x_n) \not\rightarrow f(x_0)$.

□