

## Sample Exam 2 – Version 2

*No calculators. Show your work. Give full explanations. Good luck!*

1. (15 points)

- (a) Carefully state the definition of what it means to say that  $\sum_{n=1}^{\infty} a_n$  is convergent.
- (b) Use this definition to prove that if  $\sum_{n=1}^{\infty} a_n$  is convergent and  $\sum_{n=1}^{\infty} b_n$  is divergent (not convergent), then  $\sum_{n=1}^{\infty} (a_n + b_n)$  is divergent.
- (c) Prove that if  $0 \leq a_n \leq c_n$  and  $\sum_{n=1}^{\infty} c_n$  is convergent, then  $\sum_{n=1}^{\infty} a_n$  is also convergent.

2. (15 points)

- (a) Show that if  $\lim_{n \rightarrow \infty} \sqrt{n}a_n = 2$ , then  $\sum_{n=1}^{\infty} a_n$  diverges.
- (b) Find all  $x \in \mathbb{R}$  for which  $\sum_{n=1}^{\infty} \frac{x^n}{n3^{n+1}}$  converges.
- (c) Find a sequence  $\{a_n\}$  so that  $\sum_{n=1}^{\infty} a_n x^n = \frac{4x}{2-x}$  for all  $|x| < 2$ .

3. (15 points)

- (a) Carefully state the  $\varepsilon$ - $\delta$  definition of what it means for a function  $f : \mathbb{R} \rightarrow \mathbb{R}$  to be *continuous* at a point  $x_0 \in \mathbb{R}$ . Use this to show that  $f(x) = \frac{2x+1}{x^2+1}$  is continuous at  $x_0 = 2$ .
- (b) Prove that if a function  $f : \mathbb{R} \rightarrow \mathbb{R}$  is *continuous* at  $x_0$ , then  $\lim_{n \rightarrow \infty} f(x_n) = f(x_0)$  for all sequences  $\{x_n\}$  with  $\lim_{n \rightarrow \infty} x_n = x_0$ . Use this to show that

$$g(x) = \begin{cases} \cos(x^{-2}) & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$$

is not continuous at  $x_0 = 0$ .

4. (5 points) Give examples of the following, no proofs are required:

- (a) A function that is continuous at 0 and discontinuous on  $\mathbb{R} \setminus \{0\}$ .
- (b) A series with bounded partial sums that is divergent.
- (c) Bonus Points:

A sequence  $\{b_n\}$  with  $0 \leq b_n \leq \frac{1}{n}$  for each  $n \in \mathbb{N}$ , but for which  $\sum_{n=1}^{\infty} (-1)^{n+1} b_n$  diverges.

## Math 3100 - Sample Exam 2 (Version 2) - SOLUTIONS

1. (a)  $\sum_{n=1}^{\infty} a_n$  converges  $\Leftrightarrow \lim_{n \rightarrow \infty} (a_1 + a_2 + \dots + a_n)$  exists.

(b) Claim

If  $\sum_{n=1}^{\infty} a_n$  convs &  $\sum_{n=1}^{\infty} b_n$  divs, then  $\sum_{n=1}^{\infty} (a_n + b_n)$  diverges.

Proof

Since  $\sum_{n=1}^{\infty} a_n$  conv we know  $\lim_{n \rightarrow \infty} (a_1 + \dots + a_n)$  exists.

Since  $\sum_{n=1}^{\infty} b_n$  div we know  $\lim_{n \rightarrow \infty} (b_1 + \dots + b_n)$  does not exist.

IF  $\sum_{n=1}^{\infty} (a_n + b_n)$  converged this would mean that

$$\lim_{n \rightarrow \infty} ((a_1 + b_1) + \dots + (a_n + b_n)) \text{ exists.}$$

$$\text{Since } b_1 + \dots + b_n = \underbrace{((a_1 + b_1) + \dots + (a_n + b_n))}_{\text{convergent sequence}} - \underbrace{(a_1 + \dots + a_n)}_{\text{convergent sequence}}$$

this would contradict  $\sum_{n=1}^{\infty} a_n$  conv &  $\sum_{n=1}^{\infty} b_n$  div □.

(c) Claim (Direct Comparison Test).

If  $0 \leq a_n \leq c_n$  &  $\sum_{n=1}^{\infty} c_n$  convergent, then  $\sum_{n=1}^{\infty} a_n$  converges

Proof Let  $L = \lim_{n \rightarrow \infty} (c_1 + \dots + c_n)$ . Since  $0 \leq a_n \leq c_n \forall n \in \mathbb{N}$  it follows that  $(a_1 + \dots + a_n) \leq L \forall n \in \mathbb{N}$ . Since the seq of partial sums  $\{a_1 + \dots + a_n\}$  is also increasing it follows from the MCT that  $\lim_{n \rightarrow \infty} (a_1 + \dots + a_n)$  exists  $\Leftrightarrow \sum_{n=1}^{\infty} a_n$  converges.

2. (a) Claim

Claim  
If  $\lim_{n \rightarrow \infty} \sqrt[n]{a_n} = 2$ , then  $\sum_{n=1}^{\infty} a_n$  diverges.

Proof

Since  $\sqrt[n]{a_n} = \frac{a_n}{\frac{1}{\sqrt[n]{a_n}}} \rightarrow 2$  and  $\sum_{n=1}^{\infty} \frac{1}{\sqrt[n]{a_n}}$  diverges

it follows from the "limit comparison test" that  $\sum_{n=1}^{\infty} a_n$  diverges  $\square$

(b) Claim  $\sum_{n=1}^{\infty} \frac{x^n}{n 3^{n+1}}$  converges  $\Leftrightarrow x \in [-3, 3)$

Pract Let  $a_n = \frac{x^n}{n3^{n+1}}$ .

Since  $\left| \frac{a_{n+1}}{a_n} \right| = \frac{n}{n+1} \cdot \frac{1}{3} |x| \rightarrow \frac{1}{3} |x|$  it follows from

the "Ratio Test" that  $\sum_{n=1}^{\infty} a_n$  conv. abs. if  $|x| < 3$

and diverges if  $|x| > 3$ .

If  $x=3$  then  $\sum_{n=1}^{\infty} a_n = \sum_{n=1}^{\infty} \frac{1}{3^n}$  which diverges.

If  $x = -3$  then  $\sum_{n=1}^{\infty} a_n = \sum_{n=1}^{\infty} (-1)^n \frac{1}{3^n}$  which converges

by the Alt. Series Test since  $\frac{1}{3n} \searrow 0$ .

(c) Since  $\frac{4x}{2-x} = 2 \times \frac{1}{1-\frac{x}{2}}$  &  $\frac{1}{1-\frac{x}{2}} = \sum_{n=0}^{\infty} \frac{x^n}{2^n} = \sum_{n=1}^{\infty} \frac{x^{n-1}}{2^{n-1}}$  if  $|x| < 2$

it follows that  $\frac{4x}{2-x} = \sum_{n=1}^{\infty} \left( \frac{1}{2^{n-1}} \right) x^n$  if  $|x| < 2$

3. (a) (i) See Sample Exam 2 (Version 1) Q3(a)(i).

(ii) Claim:  $f(x) = \frac{2x+1}{x^2+1}$  is continuous at  $x_0=2$ .

Proof Let  $\varepsilon > 0$  and set  $\delta = \min\{1, \frac{\varepsilon}{3}\}$ .

If  $|x-x_0| < \delta$ , then

$$\left| \frac{2x+1}{x^2+1} - 1 \right| = \frac{|x|}{x^2+1} |x-2| \leq |x| |x-2| < 3|x-2| < 3\left(\frac{\varepsilon}{3}\right) = \varepsilon$$

$\uparrow$   
 $f(2)$

since  $x^2 > 0$  always.

since  $|x-2| < \frac{\varepsilon}{3}$ .

since  $|x-2| < 1$   
 $\Rightarrow |x| < 3$

□

(b) (i) See Sample Exam 2 (Version 1) Q3(c) direction ( $\Rightarrow$ ).

(ii) Claim  $g(x) = \begin{cases} \cos(x^{-2}) & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$  is discontinuous at  $x_0=0$ .

Proof By (i) it suffices to exhibit two sequences  $\{x_n\}$  &  $\{y_n\}$  with  $\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} y_n = 0$ , but  $\lim_{n \rightarrow \infty} g(x_n) \neq \lim_{n \rightarrow \infty} g(y_n)$ .

Take  $x_n = \frac{1}{\sqrt{2\pi n}}$  and  $y_n = \frac{1}{\sqrt{2\pi n + \frac{\pi}{2}}}$  for each  $n \in \mathbb{N}$ .

Clearly  $x_n \rightarrow 0$  and  $y_n \rightarrow 0$ , while  $g(x_n) = 1$  &  $g(y_n) = 0 \forall n \in \mathbb{N}$ .

□

4. (a)  $f(x) = \begin{cases} x & \text{if } x \in \mathbb{Q} \\ 0 & \text{if } x \in \mathbb{R} \setminus \mathbb{Q} \end{cases}$ .

(b)  $\sum_{n=1}^{\infty} (-1)^n$

(c)  $b_n = \begin{cases} \frac{1}{n} & \text{if } n \text{ even} \\ 0 & \text{if } n \text{ odd} \end{cases}$ .