

Final Exam - Practice Questions

1. Evaluate the given line integral.
 - (a) $\int_C yz^2 ds$, where C is the line segment from $(-1, 1, 3)$ to $(0, 3, 5)$
 - (b) $\int_C x^3 z ds$, where $C : x = 2 \sin t, y = t, z = 2 \cos t, 0 \leq t \leq \pi/2$
 - (c) $\int_C x^3 y dx - x dy$, where C is the circle $x^2 + y^2 = 1$ with counterclockwise orientation
 - (d) $\int_C x \sin y dx + xyz dz$, where C is given by $\mathbf{r}(t) = t \mathbf{i} + t^2 \mathbf{j} + t^3 \mathbf{k}, 0 \leq t \leq 1$
 - (e) $\int_C \mathbf{F} \cdot d\mathbf{r}$, where $\mathbf{F}(x, y) = x^2 y \mathbf{i} + e^y \mathbf{j}$ and C is given by $\mathbf{r}(t) = t^2 \mathbf{i} - t^3 \mathbf{j}, 0 \leq t \leq 1$
 - (f) $\int_C \mathbf{F} \cdot d\mathbf{r}$, where $\mathbf{F}(x, y, z) = \langle x + y, z, x^2 y \rangle$ and C is given by $\mathbf{r}(t) = \langle 2t, t^2, t^4 \rangle, 0 \leq t \leq 1$
2. Find the work done by the force field $\mathbf{F}(x, y, z) = z \mathbf{i} + x \mathbf{j} + y \mathbf{k}$ in moving a particle from the point $(3, 0, 0)$ to the point $(0, \pi/2, 3)$
 - (a) along a straight line
 - (b) along the helix $x = 3 \cos t, y = t, z = 3 \sin t$
3. Show that \mathbf{F} is a conservative vector field and find a potential function f such that $\mathbf{F} = \nabla f$.
 - (a) $\mathbf{F}(x, y) = \sin y \mathbf{i} + (x \cos y + \sin y) \mathbf{j}$
 - (b) $\mathbf{F}(x, y, z) = (2xy^3 + z^2) \mathbf{i} + (3x^2 y^2 + 2yz) \mathbf{j} + (y^2 + 2xz) \mathbf{k}$
4. Show that \mathbf{F} is a conservative vector field and use this fact to evaluate $\int_C \mathbf{F} \cdot d\mathbf{r}$ along the given curve.
 - (a) $\mathbf{F}(x, y) = (2x + y^2 + 3x^2 y) \mathbf{i} + (2xy + x^3 + 3^2) \mathbf{j}$, where C is the arc of the curve $y = x \sin x$ from $(0, 0)$ to $(\pi, 0)$
 - (b) $\mathbf{F}(x, y, z) = yz(2x + y) \mathbf{i} + xz(x + 2y) \mathbf{j} + xy(x + y) \mathbf{k}$, where C is given by $\mathbf{r}(t) = (1 + t) \mathbf{i} + (1 + 2t^2) \mathbf{j} + (1 + 3t^3) \mathbf{k}, 0 \leq t \leq 1$
5. Verify that Green's Theorem is true for the line integral

$$\int_C xy dx + x^2 dy$$

where C is the triangle with vertices $(0, 0)$, $(1, 0)$, and $(1, 2)$.

6. Use Green's Theorem to evaluate

$$\int_C (1 + \tan x) dx + (x^2 + e^y) dy$$

where C is the positively oriented boundary of the region enclosed by the curves $y = \sqrt{x}$, $x = 1$, and $y = 0$.

7. Find the counterclockwise circulation and outward flux of the field $\mathbf{F}(x, y) = (-\sin x) \mathbf{i} + x \cos y \mathbf{j}$ around and over the square cut from the first quadrant by the lines $x = \pi/2$ and $y = \pi/2$.
8. Use Green's Theorem to find the work done by the force $\mathbf{F}(x, y) = x(x + y) \mathbf{i} + xy^2 \mathbf{j}$ in moving a particle from the origin along the x -axis to $(1, 0)$, then along the line segment to $(0, 1)$, and then back to the origin along the y -axis

9. Use Green's Theorem to find the area of the regions bounded by the curves with the following vector equation.

(a) $\mathbf{r}(t) = \cos^3 t \mathbf{i} + \sin^3 t \mathbf{j}$, $0 \leq t \leq 2\pi$

(b) $\mathbf{r}(t) = \cos t \mathbf{i} + \sin^3 t \mathbf{j}$, $0 \leq t \leq 2\pi$

10. Evaluate

$$\int_C y^2 dx + 3xy dy$$

where C is the positively oriented boundary of the semi-annular region R in the upper half-plane between the circles $x^2 + y^2 = 1$ and $x^2 + y^2 = 4$.

11. Find the area of the part of the surface $z = x^2 + 2y$ that lies above the triangle with vertices $(0, 0)$, $(1, 0)$, and $(1, 2)$.

12. Evaluate the given surface integral.

(a) $\iint_S z d\sigma$, where S is the part of the paraboloid $z = x^2 + y^2$ that lies under the plane $z = 4$

(b) $\iint_S (x^2 z + y^2 z) d\sigma$, where S is the part of the plane $z = 4 + x + y$ that lies inside the cylinder $x^2 + y^2 = 4$

(c) $\iint_S \mathbf{F} \cdot \mathbf{n} d\sigma$, where $\mathbf{F}(x, y, z) = xz \mathbf{i} - 2y \mathbf{j} + 3x \mathbf{k}$ and S is the sphere $x^2 + y^2 + z^2 = 4$ with outward orientation

13. Verify Stokes' Theorem is true for the vector field $\mathbf{F}(x, y, z) = y \mathbf{i} + z \mathbf{j} + x \mathbf{k}$, where S is the part of the plane $x + y + z = 1$ that lies in the first octant, orientated upwards.

14. Use Stokes' Theorem to evaluate

$$\iint_S \text{curl } \mathbf{F} \cdot \mathbf{n} d\sigma$$

where $\mathbf{F}(x, y, z) = x^2 y z \mathbf{i} + y z^2 \mathbf{j} + z^3 e^{xy} \mathbf{k}$ and S is the part of the sphere $x^2 + y^2 + z^2 = 5$ that lies above the plane $z = 1$ and is oriented upward.

15. Use Stokes' Theorem to evaluate

$$\int_C \mathbf{F} \cdot d\mathbf{r}$$

where $\mathbf{F}(x, y, z) = xy \mathbf{i} + yz \mathbf{j} + zx \mathbf{k}$ and C is the triangle with vertices $(1, 0, 0)$, $(0, 1, 0)$, and $(0, 0, 1)$, oriented counterclockwise as viewed from above.

16. Verify that the Divergence Theorem is true for the vector field $\mathbf{F}(x, y, z) = x \mathbf{i} + y \mathbf{j} + z \mathbf{k}$, where D is the unit ball $x^2 + y^2 + z^2 \leq 1$.

17. Use the Divergence Theorem to calculate the surface integral

$$\iint_S \mathbf{F} \cdot \mathbf{n} d\sigma$$

where $\mathbf{F}(x, y, z) = x^3 \mathbf{i} + y^3 \mathbf{j} + z^3 \mathbf{k}$ and S is the surface of the solid bounded by the cylinder $x^2 + y^2 = 1$ and the planes $z = 0$ and $z = 2$.

18. Compute the outward flux of

$$\mathbf{F}(x, y, z) = x^2 \mathbf{i} + y^2 \mathbf{j} + z^2 \mathbf{k}$$

through

(a) the cube cut from the first octant by the planes $x = 1$, $y = 1$, and $z = 1$

(b) the cube bounded by the planes $x = \pm 1$, $y = \pm 1$, and $z = \pm 1$

19. Find $\text{curl } \mathbf{F}$ and $\text{div } \mathbf{F}$ if $\mathbf{F}(x, y, z) = x^2 z \mathbf{i} + 2x \sin y \mathbf{j} + 2z \cos y \mathbf{k}$.

20. Show that there is no vector field \mathbf{G} such that $\text{curl } \mathbf{G} = 2x \mathbf{i} + 3yz \mathbf{j} + xz^2 \mathbf{k}$.