

# Math 8100 Assignment 7

## Hilbert Spaces

Due date: Friday 9th of November 2018

1. (a) Prove that  $\ell^2(\mathbb{N})$  is complete.

Recall that  $\ell^2(\mathbb{N}) := \{x = \{x_j\}_{j=1}^\infty : \|x\|_{\ell^2} < \infty\}$ , where  $\|x\|_{\ell^2} := \left(\sum_{j=1}^\infty |x_j|^2\right)^{1/2}$ .

- (b) Let  $H$  be a Hilbert space. Prove the so-called *polarization identity*, namely that for any  $x, y \in H$ ,

$$\langle x, y \rangle = \frac{1}{4} (\|x + y\|^2 - \|x - y\|^2 + i\|x + iy\|^2 - i\|x - iy\|^2)$$

and conclude that any invertible linear map from  $H$  to  $\ell^2(\mathbb{N})$  is *unitary* if and only if it is *isometric*.

Recall that if  $H_1$  and  $H_2$  are Hilbert spaces with inner products  $\langle \cdot, \cdot \rangle_1$  and  $\langle \cdot, \cdot \rangle_2$ , then a mapping  $U : H_1 \rightarrow H_2$  is said to be **unitary** if it is an invertible linear map that preserves inner products, namely  $\langle Ux, Uy \rangle_2 = \langle x, y \rangle_1$ , and an **isometry** if it preserves “lengths”, namely  $\|Ux\|_2 = \|x\|_1$ .

2. Let  $E$  be a subset of a Hilbert space  $H$ .

(a) Show that  $E^\perp := \{x \in H : \langle x, y \rangle = 0 \text{ for all } y \in E\}$  is a closed subspace of  $H$ .

(b) Show that  $(E^\perp)^\perp$  is the smallest closed subspace of  $H$  that contains  $E$ .

3. In  $L^2([0, 1])$  let  $e_0(x) = 1$ ,  $e_1(x) = \sqrt{3}(2x - 1)$  for all  $x \in (0, 1)$ .

(a) Show that  $e_0, e_1$  is an orthonormal system in  $L^2(0, 1)$ .

(b) Show that the polynomial of degree 1 which is closest with respect to the norm of  $L^2(0, 1)$  to the function  $f(x) = x^2$  is given by  $g(x) = x - 1/6$ . What is  $\|f - g\|_2$ ?

4. (a) Verify that the following systems are orthogonal in  $L^2([0, 1])$ :

i.  $\{1/\sqrt{2}, \cos(2\pi x), \sin(2\pi x), \dots, \cos(2\pi kx), \sin(2\pi kx), \dots\}$

ii.  $\{e^{2\pi i k x}\}_{k=-\infty}^\infty$

(b) Let  $f \in L^1([0, 1])$ .

i. Show that for any  $\epsilon > 0$  we can write  $f = g + h$ , where  $g \in L^2$  and  $\|h\|_1 < \epsilon$ .

ii. Use this decomposition of  $f$  to prove the so-called *Riemann-Lebesgue lemma*:

$$\lim_{k \rightarrow \infty} \int_0^1 f(x) \cos(2\pi kx) dx = \lim_{k \rightarrow \infty} \int_0^1 f(x) \sin(2\pi kx) dx = 0$$

5. (a) The first three Legendre polynomials are

$$P_0(x) = 1, \quad P_1(x) = x, \quad P_2(x) = (3x^2 - 1)/2.$$

Show that the orthonormal system in  $L^2([-1, 1])$  obtained by applying the Gram-Schmidt process to  $1, x, x^2$  are scalar multiples of these.

(b) Compute

$$\min_{a,b,c} \int_{-1}^1 |x^3 - a - bx - cx^2|^2 dx$$

(c) Find

$$\max \int_{-1}^1 x^3 g(x) dx$$

where  $g$  is subject to the restrictions

$$\int_{-1}^1 g(x) dx = \int_{-1}^1 xg(x) dx = \int_{-1}^1 x^2g(x) dx = 0; \quad \int_{-1}^1 |g(x)|^2 dx = 1.$$

6. Let

$$\mathcal{C} = \left\{ f \in L^2([0, 1]) : \int_0^1 f(x) dx = 1 \text{ and } \int_0^1 xf(x) dx = 2 \right\}$$

(a) Let  $g(x) = 18x^2 - 5$ . Show that  $g \in \mathcal{C}$  and that

$$\mathcal{C} = g + \mathcal{S}^\perp$$

where  $\mathcal{S}^\perp$  denotes the orthogonal complement of  $\mathcal{S} = \text{Span}(\{1, x\})$ .

(b) Find *the* function  $f_0 \in \mathcal{C}$  for which

$$\int_0^1 |f_0(x)|^2 dx = \inf_{f \in \mathcal{C}} \int_0^1 |f(x)|^2 dx.$$

### Extra Challenge Problems

*Not to be handed in with the assignment*

1. Prove that every closed convex set  $K$  in a Hilbert space has a unique element of minimal norm.

2. **The Mean Ergodic Theorem:** Let  $U$  be a unitary operator on a Hilbert space  $H$ .

*Prove that if  $M = \{x : Ux = x\}$  and  $S_N = \frac{1}{N} \sum_{n=0}^{N-1} U^n$ , then  $\lim_{N \rightarrow \infty} \|S_N x - Px\| = 0$  for all  $x \in H$ , where  $Px$  denotes the orthogonal projection of  $x$  onto  $M$ .*